



## Missouri Pharmacy Program – Preferred Drug List



### Tramadol-Like Agents

**Effective 05/05/2010**

Revised 06/21/2011

#### Preferred Agents

- Tramadol
- Tramadol-Apap
- Tramadol ER

#### Non-Preferred Agents

- Ultram®
- Ultracet®
- Ultram ER®
- Ryzolt ER®
- Nucynta®
- Rybix ODT®

<u>Approval Criteria</u>	<u>Denial Criteria</u>
Failure to achieve desired therapeutic outcomes with trial on 2 preferred agents	See Below*
Documented trial period for preferred agents	
Documented ADE/ADR to preferred agents	
Documented compliance on current therapy regimen	Drug Prior Authorization Hotline: (800) 392-8030

#### **\*Denial Criteria**

##### **Ultram:**

- 16 years of age or less<sup>2</sup>
- Daily dose exceeds maximum recommended daily dose:
  - Patients  $\leq$  75 years of age – 400 mg
  - Patients  $>$  75 years of age – 300 mg
- Presence of seizure history
- Concurrent use of monoamine-oxidase inhibitor or discontinuation of one within 14 days

**Ultracet:**

- 15 years of age or less
- Daily dose exceeds maximum recommended daily dose:
  - Patients  $\geq$  16 years of age – 300 mg tramadol and 2600mg acetaminophen
- Presence of seizure history
- Concurrent use of monoamine-oxidase inhibitor or discontinuation of one within 14 days

**Ultram ER/Ryzolt:**

- 15 years of age or less
- Daily dose exceeds maximum recommended daily dose:
  - Patients  $\geq$  16 years of age – 300 mg
- Presence of seizure history
- Concurrent use of monoamine-oxidase inhibitor or discontinuation of one within 14 days
- History of opioid abuse or dependence – therapy will be subject to Clinical Consultant review
- Claim for Ultram ER/Ryzolt therapy without previous stabilization on immediate-release tramadol product.

**Nucynta:**

- 18 years of age or less
- Daily dose exceeds maximum recommended daily dose:
  - Patients  $\geq$  18 years of age – 700 mg
- Presence of seizure history
- Concurrent use of monoamine-oxidase inhibitor or discontinuation of one within 14 days

**Rybix ODT**

- 16 years of age or less
- Daily dose exceeds maximum recommended daily dose:
  - Patients  $\geq$  17 years of age – 400mg
- Presence of seizure history
- Concurrent use of monoamine-oxidase inhibitor or discontinuation of one within 14 days